

Bekenstein-Hawking entropy from a phenomenological membrane

Carsten Gundlach

LAEFF-INTA, PO Box 50727, 28080 Madrid, Spain

(9 January 1995)

It is pointed out that the entropy of a membrane which is quantized perturbatively around a background position of fixed radius in a black hole spacetime is equal to the Bekenstein-Hawking entropy, if 1) the membrane surface is the horizon surface plus one Planck unit, and 2) its temperature is the Hawking temperature. (This is a comment on gr-qc 9411037.)

This note is a comment on the recent preprint [3], where a Nambu-Goto membrane was proposed as the site of black hole entropy, but where the resulting quantum theory was subsequently confused with that of a single point particle on the two-sphere $r = \text{const.}$, resulting in an entropy of order $\ln(M)$. We now give the correct calculation. Consider a massless membrane with Nambu-Goto action (volume of its worldsheet) in the background spacetime of a Schwarzschild or Reissner-Nordström black hole. With r and t the usual Schwarzschild radial and time coordinates and θ and ϕ the usual polar coordinates in spherical symmetry, let the membrane worldsheet be parametrized by $r(t, \theta, \phi) = R(t) + \eta(t, \theta, \phi)$. The zero mode $R(t)$ is determined by imposing that $\int \eta \sin \theta d\theta d\phi = 0$ for all t . We decompose $\eta(t, \theta, \phi)$ into a spherical harmonics as $\sum_{l,m} \eta_{lm}(t) Y_{lm}(\theta, \phi)$. Expanding to quadratic order in the η_{lm} we obtain a background action plus a system of harmonic oscillators η_{lm} parametrically coupled to the zero mode R . Now we make the assumption that $R(t)$ is fixed at a constant value $\bar{R}$. This is of course inconsistent, as there is no such background solution other than $\bar{R}$ equal to the horizon radius. We quantize the η_{lm} around this background value. The energy levels for each independent harmonic oscillator η_{lm} are given by

$$E_{lm} = \left(N_{lm} + \frac{1}{2} \right) [E_0^2(\bar{R}) + E_1^2(\bar{R})l(l+1)]^{1/2}, \quad (1)$$

where $E_1^2(r) = (r - r_-)(r - r_+)r^4$ for a Reissner-Nordström black hole, or $E_1^2(r) = (r - 2M)/r^3$ for a Schwarzschild black hole. The N_{lm} for each (l, m) , where $l = 1, 2, \dots$ and $m = -l, \dots, l$, are the quantum numbers for the system. In the partition function we sum over all values $N_{lm} = 0, 1, 2, \dots$ for each (l, m) . In calculating the partition function approximately, we expand for large l and N , so that $E_{lm} \sim N_{lm} E_1(\bar{R})$, and approximate a sum over l by an integral. (A similar calculation arises in t'Hooft's "brick wall" model [1].) We obtain for the free energy $F = -T \ln Z = -2E_1^{-2}(\bar{R})T^3$ and hence for the entropy $S = -\partial F / \partial T$

$$S = 6E_1^{-2}(\bar{R})T^2 \quad (2)$$

So far this is only quantum mechanics, and Newton's constant has not yet appeared. To this perturbative level, the energy levels and hence the entropy are also independent of the membrane tension. We now parametrize $\bar{R}$ as $\bar{R} = 2M + \delta/(16\pi M)$ (now in Planck units $G = 1$), which means that its area is δ Planck units larger than the horizon area. It has been proposed that $\delta \sim 1$ for a phenomenological membrane ("stretched horizon"). We obtain $E_1 = (1/4M^2)\sqrt{\delta/4\pi}$. Assuming that the temperature in our entropy formula is the Hawking temperature, $T = 1/(8\pi M)$, we obtain for the entropy

$$S = 6M^2/(\pi\delta) \quad (3)$$

We see that a δ of order 1 gives the Bekenstein-Hawking entropy.

[1] G. t'Hooft, Nucl. Phys. **B256**, 727 (1985).

[2] L. Susskind, L. Thorlacius, and J. Uglum, Phys. Rev. D **48**, 3743 (1993).

[3] C. O. Lousto and M. Maggiore, On the energy spectrum of quantum black holes, preprint gr-qc 9411037.